

BF

(4)

19  **Europäisches Patentamt**
European Patent Office
Office européen des brevets

11 Publication number: **0 179 441**
A2

12 **EUROPEAN PATENT APPLICATION**

21 Application number: 85113383.5

51 Int. Cl.⁴: **A 23 L 1/03**
A 21 D 2/30

22 Date of filing: 22.10.85

30 Priority: 23.10.84 SE 8405295

43 Date of publication of application:
30.04.86 Bulletin 86/18

84 Designated Contracting States:
AT BE CH DE FR GB IT LI LU NL SE

71 Applicant: Sirén, Matti
Casa Camboni Via al Crespo
CH-6596 Gordola(CH)

72 Inventor: Sirén, Matti
Casa Camboni Via al Crespo
CH-6596 Gordola(CH)

74 Representative: Eitle, Werner, Dipl.-Ing. et al,
Hoffmann, Eitle & Partner Patentanwälte Arabellastrasse
4
D-8000 München 81(DE)

54 A method of making a food composition containing inositoltriphosphate and the composition.

57 A method of making a food composition and the composition containing at least 5 mg of inositoltriphosphate.

EP 0 179 441 A2

A METHOD OF MAKING A FOOD COMPOSITION CONTAINING
INOSITOLTRIPHOSPHATE AND THE COMPOSITION

FIELD OF INVENTION

The present invention relates to a method of making an improved foodcomposition containing at least 5 mg inositoltriphosphate (IP_3) per 100 g composition and such composition having such content of IP_3 .

There is an increasing need for counteracting any bad influence of civilization, for instance environmental dangers caused by input of dangerous materials like heavy metals and radiation.

There is also a need for counteracting the hazards of smoking and other bad habits by development of healthy diet and nutrition agents.

Even as early as the year 1900, different researchers had reported the finding of the organic phosphate compound phytic acid, i.e., 1,2,3,4,5,6-hexakis (dihydrogenphosphate) myo-inositol (also sometimes called inositol-hexaphosphoric acid) in plants. The content of phytic acid in different plants varies considerably. The content in grain is usually approximately 0.5-2%, with certain exceptions. Polished rice has a level of only 0.1% while wild rice contains as much 2.2% phytic acid. Beans contain about 0.4-2%, oil plants approximately 2-5% and pollen 0.3-2%. The content of phytic acid in the plant varies during the growth period. The content is also influenced by, among other things, the climate.

In the literature there are reports on the presence of inositol pentaphosphate (IP_5) and inositol tetraphosphate (IP_4) in a few plants. It is further known that phosphate derivatives lower than IP_6 are formed at germination of grain. For instance the final products at the germination are inositol and phosphate. The use of IP_6 has been described in several scientific publications. The majority of the authors of these articles have observed several negative effects on humans and animals when consuming IP_6 or substances containing IP_6 . Feeding dogs with too high an amount of IP_6 gives rise for example to rachitis. In humans lack of zinc and as a consequence thereof slower growth of children has been observed. Anemia has been observed mainly in women. Because of the above mentioned negative effects on the mineral balance in humans and animals, attempts have so far been made to reduce the intake of IP_6 and its derivatives to a minimum.

Cadmium also has been found to be detrimental to human health. While this metal in general is present in a low level in our environment, the amount of cadmium we are exposed to depends on several factors. Cadmium occurrence as well as its availability in the ground varies among different areas, with a relatively high uptake in plants growing in areas with relatively low pH value. By industrial activity, mainly handling of metals, cadmium can be released into the air, ground and water. Cadmium in soil is absorbed by plants and thus can come into the diet of human beings and animals. The most important routes of exposure to cadmium are via smoking, food and, to a certain extent, drinking water.

Cadmium is mainly absorbed in the intestine and through the lungs, although only a small part of the cadmium in the diet is absorbed. The average cadmium intake via food is estimated to be approximately 50 ug per day in most countries, but the variation is large among different geographic areas and among individuals. Data from smokers show that as much as 50% of the inhaled cadmium can be absorbed. Several investigations show twice as high blood- and organ-levels of cadmium in smokers compared to non-smokers. The excretion of cadmium from the human body is slow and a half-life of 10-30 years has been reported. This means that cadmium is accumulated in the body. The main part, 80-90% of the accumulated cadmium, is bound to a protein, metallothionein, mainly in the liver and kidneys. The formation of metallothionein is induced by metals, mainly zinc and cadmium. The binding of cadmium to metallothionein is very strong and results in a detoxification of cadmium. The remaining cadmium is in the body, i.e. that not bound to metallothioneins, is distributed among the other organs of the body with relatively high levels in the intestine, lungs (especially of smokers), the circulatory system (heart, artery walls, spleen) and glands like the pancreas and prostate.

Among the negative effects, it is known that cadmium can affect the elastin/elastase system of the body. It is also known that cadmium can affect several different enzymes in the body, examples of which are Na^+ , K^+ (Mg^{2+})-ATP-ase and Ca^{2+} , Mg^{2+} -ATP-ase, which are important in ion transport systems. Further examples are cytochrome-P450-enzymes which hydrolyze steroids, fatty acids, aromatic compounds and toxic compounds. Other important enzymes, which are inhibited by cadmium, are glutathion-peroxidase, and superoxiddismutase, which protect against occurrence of peroxidation. Zinc dependent enzymes, such as leucine-aminopeptidase, are also inhibited by cadmium.

Results from a large number of animal experiments obtained over many years show negative effects even at very low levels of cadmium. This would mean that a large proportion of the population is negatively affected, and this is above all valid for smokers. Epidemiological research shows a connection between the presence of cancer, high blood pressure and cardiovascular diseases (for instance, arteriosclerosis, heart infarction, sudden heart death) and the occurrence of cadmium in the environment. Exposure to cadmium also seems to be a factor in increasing the risk of age diabetes.

There are also investigations showing that cadmium can have negative effects on the kidneys, lungs (fibrosis, emphysema, cancer), blood vessel walls (fat deposition, arteriosclerosis, vessel wall contraction, elasticity, damage to endothelium), prostacycline production, prostate, heart (conduction system, force of contraction), placenta, testicles and central nerve system. Cadmium can also induce the formation of free radicals and thereby cause lipid peroxidation, which can be important in the origin of other diseases like rheumatism. Allergies and bronchitis can also be connected with cadmium exposure. The knowledge of the negative influence of cadmium on humans and animals has increased considerably over the last decades.

0179441

A very intensive research effort has been made for many years seeking to counteract the above mentioned negative effects of heavy metals, such as cadmium and the negative effects of free radicals which are formed in different ways, for instance by metals such as iron, aluminium and cadmium and by radiation. Of course, also the hazards of smoking have been studied for a long time.

SUMMARY OF THE INVENTION

According to the present invention it has been possible to avoid or at least alleviate the above negative effects observed on humans and animals, by consumption of the special inositolphosphate IP_3 . Thus, the invention provides a method of making a food composition and the said composition containing at least 5 mg of IP_3 per 100 g of composition. Generally, it is preferred to use the IP_3 in salt form. However, it can also be used in acid form, if desired.

DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

According to the present invention a food composition having a content of IP_3 of at least 5 mg per 100 g composition has been brought about. Very often the IP_3 content is at least 20 mg per 100 g composition. Advantageously, the content of IP_3 should be within the range of 5-500, preferably in the range of 20-500, more preferably 50-500, 100-500 or 150-500 mg IP_3 per 100 g food composition.

0179441

The composition can be used as an additive or an intermediate concentrate to increase the IP_3 content of other foodstuff products. Then the content of IP_3 in said intermediate concentrate should be at least 20 mg per 100 g of the concentrate. Usually, however, the content of IP_3 in the concentrate is much higher, advantageously 50 mg - 100 g and with preference 75 mg - 80 g, 100 mg - 80 g, 150 mg - 60 g, 200 mg - 60 g, 250 mg - 50 g or 300 mg - 50 g respectively per 100 g concentrate. Preferably, the content of IP_3 of the concentrate should be as high as possible.

The intermediate concentrate can be used in many different forms, such as powder, tablets, capsules and granules. However, it is also possible to use it in the form of a liquid, such as an aqueous solution.

The IP_3 is preferably selected from the group consisting of D-myo-inositol-1.2.6-triphosphate, D-myo-inositol-1.2.5-triphosphate, myo-inositol-1.2.3-triphosphate, L-myo-inositol-1.3.4-triphosphate and D-myo-inositol-1.4.5-triphosphate and mixtures thereof. Of these isomers D-myo-inositol-1.2.6-triphosphate is preferred.

Often 20-100, preferably 40-100 % by weight of the IP_3 content consists of D-myo-inositol-1.2.6-triphosphate.

According to one suitable method for the production of IP_3 a material containing IP_6 is broken down enzymatically with phytase enzyme. The IP_6 can be provided either as pure material or in the form of an IP_6 containing source, such as wheat bran. Phytase enzyme can be found for instance in plants, seeds and micro-organisms.

0179441

By the enzymatic treatment of the IP_6 a hydrolysis takes place resulting in a mixture of different lower inositolphosphates, i.e. inositolpentaphosphate (IP_5), inosoltetraphosphate (IP_4), inosolttriphosphate (IP_3), inosoltldiphosphate (IP_2) and inosoltmonophosphate (IP_1).

Usually, the hydrolysis is carried out at a temperature of 20-70°C and a pH of 4 to 8. The hydrolysis is suitable stopped when the liberation of about 30-60% of the total ester phosphorus has been achieved. At said stage a high porportion of the desired IP_3 isomer of isomers has been formed by hydrolysis of the IP_6 containing material.

The mixture of inositolphosphates obtained may hereafter be separated by chromatography to isolate the IP_3 -containing fraction. Preferably, this is made in a column. If the IP_3 fraction contains more than one isomer, these isomers are separated in another subsequent chromatographic separation step.

The IP_3 can be obtained as a salt or as an acid thereof. The salt form is preferred, since it is easier to produce in pure and concentrated form than the acid.

The salt form of the IP_3 isomer is readily obtainable from the acid form using standard procedures. Thus, there can be prepared salts, such as alkali metal and alkaline earth metal salts, e.g. lithium, sodium, potassium, calcium or magnesium. However, also the zinc salts are very useful as well as the NH_4^+ and organic amine salts. Exemplary amines are triethanolamine, diethanolamine, triisopropanolamine, N,N-dimethyl-2-amino-2-methyl-1-propanol, N,N-dimethylethanolamine, tetrabutylamine and cyclohexylamine. Also other salts might be used. Especially preferred salts are those which are physiologically acceptable.

Advantageously, the distribution curve showing the content of the different inositolphosphates has a maximum and preferably the sole maximum for IP_3 which means that the content of IP_3 is larger than IP_2 and/or IP_4 . Usually the proportion of IP_3 is at least 10 % of the total amount of inositolphosphates.

Sometimes the composition in addition to IP_3 has a content of IP_4 and/or inositoldiphosphate (IP_2). Then, preferably more than 40 % by weight of the total amount of inositolphosphates in the composition consists of IP_3 , while 30-85 % by weight of the remaining inositolphosphates consists of IP_2 plus IP_4 . In such a composition the IP_3 can consist essentially of D-myo-inositol-1.2.6-triphosphate. However, also other IP_3 isomers, especially those mentioned above, can be used in such a foodstuff product.

Depending for instance on the form of the composition the IP_3 can be present in salt form or in acid form. The acid form is usually used as a liquid, preferably an aqueous solution. In salt form the IP_3 can be used as a dry product or alternatively as a liquid, preferably an aqueous solution.

When IP_3 is present as a salt, said salt is generally selected from the group mentioned above.

The invention provides a method of making a food composition, said food being initially substantially free of IP_3 . The method comprises adding a source of IP_3 to the composition in an amount sufficient to provide a final concentration of at least 5 mg IP_3 per 100 g of composition.

The source of IP_3 can be IP_3 as such produced separately or, alternatively, be IP_6 , IP_5 and/or IP_4 in the presence of phytase for enzymatic production of IP_3 .

0179441

The term "initially substantially free of IP_3 " is intended to mean that the food composition produced in the conventional way does not contain a substantial amount of IP_3 . Thus, the content of IP_3 will be less than 5 mg, normally less than 2 mg and most often below 1 mg per 100 g of the composition.

The invention also comprises a method of making a food composition wherein in the materials composing the composition, a content of phytase and of an inositolphosphate selected from the group consisting of inositoltetraphosphate (IP_4), inositolpentaphosphate (IP_5) and inositolhexaphosphate (IP_6), is established, and at at least one stage of the production process, the time and the temperature of the processing as well as the pH-value are controlled to allow an incubation in such a way that a content of inositoltriphosphate (IP_3) of at least 20 mg per 100 g composition is obtained.

At said incubation step, at least a portion of the inositolphosphate selected from IP_6 , IP_5 and IP_4 is enzymatically broken down to IP_3 with phytase enzyme. The proportion of the original inositolphosphate content transformed to IP_3 can be regulated within wide limits by varying the production parameters, such as incubation time, temperature and pH as mentioned.

Phytase enzyme may be present in plants or seeds provided they have a content of inositolhexaphosphate. Because of this it may according to the invention, not be necessary in all cases to add the enzyme if a plant or seed product is used as starting material. In the cases where said natural product has too low and enzymatic activity or when IP_6 , IP_5 or IP_4 or a mixture of these is used as starting material, a phytase enzyme, for example, from bran can be added.

0179441

Yeast can be used advantageously as a source of phytase. Preferably baker's yeast is used.

Swedish baker's yeast produced by Jästbolaget, Sweden, as well as baker's yeast produced by Rajamäki, Finland and Hefefabriken AG, Switzerland have for instance been used according to the present invention. When using yeast according to the present invention it has been established very surprisingly that essentially only one isomer is obtained, namely D-myo-inositol-1.2.6-triphosphate. Of course, the use of yeast is a very valuable procedure if said isomer only is desirable.

During the incubation a hydrolysis takes place at a suitable temperature, usually 20-70°C, preferably 30-60°C, and at a pH of 4-8. In order to stop the hydrolysis at the intended level the enzyme may be destroyed or inactivated, for instance by a rapid heating of the hydrolyzed starting material.

The method according to the invention can be modified in different ways, for instance depending on the starting material chosen. The starting material can for instance:

1. have a certain content of IP_6 , IP_5 and/or IP_4 .
2. have no content of IP_6 , IP_5 or IP_4 .

At the first alternative above, there are different possibilities to achieve a desired amount of IP_3 in the final foodstuff product. For instance the above method of hydrolyzing the inositolophosphates to IP_3 by means of phytase can be used.

0179441

If the content of IP_6 , IP_5 and/or IP_4 is not high enough in the starting material, an addition thereof can be made. In this way the IP_3 content of the final product can be increased.

The above method can be used also for the production of an intermediate product with a desired content of IP_3 , which product can be added to the starting material for the food composition.

The intermediate product can also be introduced at a later stage of the production of the foodstuff product.

Methods of producing IP_3 and its isomers as such are disclosed in applicants patent application filed on the same day as this application and having the title "Inositoltriphosphate, a method for preparing same and a composition containing same"(applicant's case No. 355b, 42762), copy of which is enclosed.

At the second alternative above, where the starting material has no content of IP_6 , IP_5 or IP_4 , such an inositolphosphate can be added together with phytase, if phytase is lacking. Then the above hydrolysis method can be used again to give the desired content of IP_3 in the final product.

Alternatively, the intermediate product mentioned above can be added to the starting material or at a later stage of the production of the food composition.

At both aforementioned methods, IP_3 in concentrated form can be added at a later or final stage of the production of the composition.

In another embodiment of this invention a method of making a food composition is provided in which the composition is initially containing less than **0179441** about 10 mg of IP_3 per 100 g of composition, wherein the content of IP_3 is increased to at least 20 mg per 100 g of composition by addition of IP_3 or a source thereof or conversion of initially contained inositolphosphate selected from the group consisting of IP_6 , IP_5 and IP_4 by enzymatic process.

In such method of making a composition according to the present invention, said composition can advantageously be an cereal based material, such as breakfast cereals, cakes, biscuits and bread. The composition can also be selected from the group consisting of sweets, chocolates and chewing gums. Often, quite preferably, the composition is also a vegetable, fruit, beverage, soup or a product based on milk, e.g. yoghurt.

In another embodiment of the invention where the composition initially is containing less than about 20 mg of IP_3 per 100 g composition. The content of IP_3 is increased to at least 50 mg per 100 g composition in the same way. In a further embodiment of the invention where the composition initially is containing less than about 50 mg of IP_3 per 100 g composition. The content of IP_3 is increased to at least 100 mg per 100 g composition in the same way.

The content of IP_3 in the composition can be varied within wide limits. It is preferred to have a content of IP_3 of 20-500, such as 50-500, 100-500 or 150-500 mg IP_3 respectively based on 100 g composition. For bakery food the interval is advantageously 20-500, preferably 100-500, and most preferably 150-500 or 200-500 mg IP_3 respectively per 100 g dry bakery food. The daily intake of IP_3 is at least 10 mg, preferably at least 50 mg.

At the production of a liquid composition according to the present invention, the content of IP_3 can also be varied to a large extent. Generally, the content of IP_3 is 5-500, such as 10-500, 20-500 or 5-300 mg IP_3 per 100 g of the liquid composition.

The liquid composition can for instance be a beverage
or soup.


0179441

According to the invention a food composition can
be provided where in the concentration of IP_3 is
at least 20 mg per 100 g of dry composition. The
 IP_3 has been added to the composition and/or formed
by adding phytase and an inositol phosphate selected
from the group consisting of IP_4 , IP_5 and IP_6 .

In the frame of the invention, bakery products are
especially preferred. Thus, a bakery food composition
is provided wherein the concentration of IP_3 is at
least 20 mg per 100 g of dry composition and the
degree of hydrolysis of naturally contained IP_6 and/or
of added IP_6 is 20-90 % preferably between 40 and
70%. Thus, the ration of inorganic phosphorus to
the total amount of phosphorus is at least 20 %,
preferably at least 40 %. At at special bakery food
composition the initial content of IP_3 is less than
150 mg per 100 g composition and the initial content
of IP_6 is less than 200 mg per 100 g composition.
The content of IP_3 in the final composition is then
increased by breaking down IP_6 .

The IP_3 -isomers mentioned above have the following
formulas:


D-myo-inositol-1.2.6-triphosphate of the formula.



where X is hydrogen, at least one univalent, divalent
or multivalent cation, or a mixture thereof, n is
the number of ions, and z is the charge of the respectively
ion;


0179441

D-myo-inositol-1.2.5-triphosphate of the formula




where X, n and z have the above mentioned meaning;


10 myo-inositol-1.2.3-triphosphate of the formula



where X, n and z have the above mentioned meaning;
L-myo-inositol-1.3.4-triphosphate of the formula



25 where X, n and z have the above mentioned meaning;
and D-myo-inositol-1.4.5-triphosphate of the formula



where X, n and z have the above mentioned meaning.

35

In each of the above formulas n ranges between 6 to 1 inclusively and z ranges from 1 to 6 inclusively. Preferably, n is between 3 to 6 inclusive and z is 3, 2 or 1.

0179441

The composition according to the present invention has a good influence on the organism in many ways. However, it is mainly intended to prevent or alleviate conditions created, induced or furthered by heavy metals, especially cadmium or diseases related to such heavy metals.

Also the composition is intended to give a good effect on smokers.

10

As examples of conditions which the present composition is intended to prevent or alleviate the following can be mentioned; high blood pressure, a cardiovascular disease, emphysema and increased platelet aggregation. However, the composition has a good effect on many other conditions too.

15

20

The invention is further explained below in connection with embodiment examples, of which examples 1 and 2 relate to a comparison test where an analysis of inositolphosphates in some commercially available breads and breakfast cereals respectively is made. Examples 3-7 illustrate a method of making bread according to the invention. Example 8 relates to the production of a cake baked on wheat flour with the addition of a calciumsalt of inositolphosphates. Example 9 shows the production of breakfast cereals after addition of a sodiumsalt of inositolphosphates. Example 10 illustrates the production of table-salt by addition of the sodiumsalt of D-myo-inositol-1.2.6-triphosphate.

25

30

35

Example 11 illustrates the production of beverages by addition of a sodiumsalt of inositoltriphosphates. Example 12 relates to the production of honey by addition of a sodiumsalt of D-myo-inositol-1.2.6-triphosphate. Example 13 illustrates the production of chocolate by addition of a sodiumsalt of inositolphosphates. Example 14 shows that in blood of rabbits,

40

platelet aggregation caused by an injection of cadmium
can be prevented by administration of a diet containing
IP₃. Example 15 shows the effect of IP₃ on the cadmium
content in different organs of mice which had got
an injection of cadmium. Example 16 shows that IP₃
prevents an increase of platelet aggregation caused
by smoking. Examples 17 and 18 show that IP₃ prevents
or reduces the formation of free radicals. Examples
19-24 illustrate hydrolysis of phytic acid in different
foodstuff sources. Examples 25-31 show production
of IP₃ and the separation thereof into different
isomers.

0179441

Example 1

Analysis of inositolphosphates in some commercially
5 available breads.

Three commercially available breads, one white bread
and two crisp breads, were analyzed for the contents
10 of inositolphosphates with HPLC. The white breads were
baked on whole rye flour and whole wheat flour respec-
tively.

15 A 20 gram quantity of the breads were ground and extracted
with 1 % hydrochloric acid for two hours at shaking.
The suspension was centrifuged and the supernatant was
collected.

20 The supernatant was analyzed with well-defined inositol-
phosphates and the results were quantified as mg inositol-
phosphates per 100 g (solid contents).

25	Type of bread	<u>IP₂</u>	<u>IP₃</u>	<u>IP₄</u>	<u>IP₅</u>	<u>IP₆</u>	(mg/100g solid content)
	White bread	33	2	3	2	2	
	Crisp bread/whole wheat flour	37	6	14	71	515	
30	Crisp bread/whole rye flour.	27	10	13	25	64	

The results show that the amount of inositoltriphosphates
35 in the commercially available breads are low.

Example 2

Analysis of inositolphosphates in breakfast cereals.

5

Commercially available Corn Flakes [®], Kellogg's was analyzed for the content of inositolphosphates with HPLC. The extraction procedure and analysis were the same as
10 described in Example 1.

19 mg IP_2 and 2 mg IP_3 per 100 g solid content was found. No IP_4 , IP_5 or IP_6 could be detected. The IP_6 contained
15 in the raw material had almost completely been broken down and the amount of IP_3 was very low.

20

25

30

35

Example 3

Variation of the fermentation period for crispbread baked
5 on rye flour.

Biologically acidified crisp bread was baked on rye flour
(1 % IP_6 content). The dough formulation was: 54.6 g
10 flour, 41.8 g water, 1.3 g salt (NaCl) and 2.4 g of a
dough from a preceding doughformulation.

A sour dough consisting of the above 2.4 g of dough from
15 a preceding doughformulation and 40 % of the flour and
85 % of the water amount was fermented in a first step
for 6 hrs before mixing with the other ingredients (flour,
water and salt). After mixing, the dough was fermented
in a second step before bread forming and baking. The
20 oven temperature was 250°C. Three breads with three dif-
ferent times for the second fermentation period were
produced. The breads were ground, extracted and analyzed
as described in Example 1. The content of IP_3 versus
fermentation time was determined as follows:

25

Second fermen- tation time	Amount of IP_3 (mg/100g solid content)
30 min	104
90 min	89
225 min	78

30

35 The result shows that an increased second fermentation
period resulted in a decrease of IP_3 content.

Example 4

IP₃-content in a bread baked on wheat and oat flour.

5

Chemically acidfied bread was baked on a combination of wheat and oat flour (0.9 % IP₆ content).

10 The dough formulation was: 37.7 % wheat flour, 17.7 % oat flour, 39.5 % water, 1.3 % salt (50 % NaCl and 50 % kalciumacetate). 1 % sucrose and 2.8 % baker's yeast.

15 After mixing the ingredients the dough was fermented before bread forming and baking. The fermentation period was 90 minutes and the temperature was 37°C. The breads obtained were ground, extracted and analyzed as described in Example 1. The content of IP₃ was determined to be
20 120 mg per 100 g dry bread.

25

30

35

Example 5

IP₃-content in a bread baked on whole wheat flour.

5

A white bread was baked on whole wheat flour (0.9 % IP₆ content). The dough formulation was: 55.9 % flour, 38.4 % water, 3.5 % yeast, 0.6 % salt (NaCl) and 1.7 % sucrose.

10

After mixing the ingredients, the dough was fermented before the bread was formed. After an additional fermentation period the bread was baked. The total fermentation period was 60 minutes and the baking temperature was 175°C.

The bread was ground, extracted and analyzed as described in Example 1. The content of IP₃ was 70 mg per 100 g dry bread.

25

30

35

0179441

Example 6

IP₃-content in a bread baked on rye flour with addition
5 of sodiumphytate.

Biologically acidified crisp bread was baked on rye flour
(1 % IP₆-content) as described in Example 3 but with
10 the difference that 0.8 g sodiumphytate was added to
100 g dough. The fermentation period was 225 minutes.

The bread was ground, extracted and analyzed as described
15 in Example 1. The amount of IP₃ in the bread was 180 mg
per 100 g dry bread.

20

25

30

35

Example 7

IP₃-content in a bread baked on rye flour with addition
5 of calciummagnesiumphytate.

Biologically acidified crisp bread was baked on rye flour
(1 % IP₆-content) as described in Example 3 but with
10 the difference that 1.5 g calciummagnesiumphytate was
added to 100 g dough. The fermentation period was 225
minutes. The bread was ground, extracted and analyzed
as described in Example 1. The amount of IP₃ in the
bread was 250 mg per 100 g dry bread.

15

20

25

30

35

Example 8

IP₃-content in a cake baked on wheat flour with the
5 addition of a calciumsalt of inositolphosphates.

A cake was baked on wheat flour (0.2 % IP₆-content).
The dough formulation was 60.1 % wheat flour, 35.7 %
10 water, 0.6 % salt (NaCl) and 3.6 % yeast.

After mixing the ingredients the dough was fermented.
0.2 g of a calciumsalt of inositolphosphates containing
15 30 % by weight of IP₃ was added in 10 ml water per 100
g dough before the cake was formed. After an additional
fermentation period the cake was baked in the oven. The
total fermentation time was 75 minutes and the baking
temperature was 225°C.

20

The cake was ground, extracted and analyzed as described
in Example 1. The amount of IP₃ was 60 mg per 100 g dry
cake.

25

30

35

Example 9

IP₃-content in breakfast cereals after addition of a sodiumsalt of inositolphosphates.

1000 g commercially available Corn Flakes ®, Kellogg's was sprayed with 10 ml warm (80°C) aqueous solution containing 50 % sucrose and 10 % of a sodiumsalt of inositolphosphates (containing 30 % IP₃). After drying, the breakfast cereals were granned, extracted and analyzed as described in Example 1. The content of IP₃ was 30 mg per 100 g dry material.

0179441

Example 10

Table-salt with addition of sodiuminositoltriphosphate.

5

The sodiumsalt of D-myo-inositol-1.2.6-triphosphate was mixed with table-salt in such a way that a final concentration of 200 ppm IP_3 was obtained.

10

15

20

25

30

35

Example 11

Beverages with addition of a sodiumsalt of inositolphosphates.

20 ml of a 15. % aqueous solution of a sodiumsalt of inositolphosphates (containing 40 % of IP_3) was added to 5 l commercially available Coca-cola[®], Seven-Up[®] and orange juice respectively. The final concentration of IP_3 in the beverages was found by HPLC to be 190 mg/l.

15

20

25

30

0179441

Example 12

Honey with addition of a sodiumsalt of inositoltriphos-
5 phates.

5 ml of a 20 % aqueous solution of the sodiumsalt of
D-myo-inositol-1.2.6-triphosphate was added to 50 kg
10 commercially available honey. The final concentration
of inositoltriphosphate was determined by HPLC to be
20 mg/kg.

15

20

25

30

35

0179441

Example 13

Chocolate with added IP_3 .

5

To 1500 g melted chocolate, 6 ml of a 10 % aqueous solution of a sodiumsalt of inositolphosphates (containing 30 % IP_3) was added before lowering the temperature and forming the final chocolate product. The content of IP_3 in the chocolate was 110 mg per kg chocolate as determined by HPLC.

15

20

25

30

35

Example 14

5 Rabbits (New Zealand white, males) weighing 2-2.5 kg were used. They were fed with a diet free from inositol phosphates, for 10 days before the experiment.

10 2 hours before the start of the experiment, 50 mg of a sodiumsalt of myo-inositol-1.2.6-triphosphate was mixed into 5 g of the diet for a group of 18 animals. Another group with 12 animals got no addition of inositoltri-phosphate.

15

Time:	Treatment
-------	-----------

20

0 minute:	Blood sample 1 (9 ml + 1 ml 3.8 % sodium citrate) taken.
-----------	--

25

1 minute:	Intravenous injection of 4 microgram Cd as CdCl_2 in 0.5 ml physiological saline, or 0.5 ml physiological saline respectively.
-----------	---

30

4 minutes:	Blood sample 2 (9 ml + 1 ml 3.8 % sodium citrate) taken.
------------	--

35

0179441

Treatment of samples

The two blood samples from each animal were centrifuged at 1200 revolutions per minute, for 10 minutes, and the plasma with platelets was obtained.

The plasma with platelets from the two samples was analyzed concerning the response to addition of ADP (adenosin diphosphate) in an aggregometer (Chronopar Corp Mod, 440) according to Born (J. Physiol: 67, 1968). The two samples were analyzed simultaneously at the same concentration (1-20 micromolar) of ADP, in the two channels of the instrument.

15

The principle of this test is that the plasma with platelets is turbid, and has a low transmittance for light. As ADP is added, the platelets aggregate, and form clumps. This results in an increase fo transmittance which is quantified by the instrument. The response to ADP was measured in scale units, with 80 scale units representing maximal aggregation. In order to have a maximal sensitivity of the method to pick up changes in platelet reactivity, the ADP dose should cause a response of 5-30 scale units. This was normally achieved with 5 uM ADP, but in some animals a lower or higher dose (1-20 uM) was necessary.

30

The result of the test is expressed as maximal aggregation in sample 2 (scale unite) minus maximal aggregation in sample 1.

35

0179441

The following results were obtained:

	<u>Oral administration</u>	<u>Injection</u>	<u>No</u>	<u>Change in aggre- gation from sample 1 to sample 2 (scale unit)</u>
5	No addition to the diet	Cd	12	+ 2.3
10	IP ₃ added to the diet	Cd	18	- 0.2

At the dose used in this experiment, the IP₃ prevented the effect of Cd on platelet aggregation.

15

An increase in platelet aggregation is regarded as one of the most important factors causing cardiovascular diseases e.g. arteriosclerosis, and the ability of IP₃ to prevent the aggregation induced by cadmium shows that IP₃ is very useful in preventing or allievating such disease.

25

30

35

Example 15

5 Mice weighing 18-20 gram at the start of the experiment were used. During the experiment and for at least seven days before the experiment the mice were fed a semisynthetic diet free of inositol phosphates. The mice were divided in two groups.

10 They received daily intraperitoneal injections of physiological saline and inositoltriphosphate (IP_3) respectively for 9 days. The dose fo IP_3 was 5.0 mg/day. The injected volume was 0.2 ml.

15 On day two of the experiment, 5-10 minutes after the second intraperitoneal injection, all mice received an intravenous injection of 2.5 microcurie of ^{109}Cd as
20 cadmium chloride in 50 ul of saline. After the last intraperitoneal injection the mice were killed and several organs were dissected out and weighed.

25 Radioactivity in the different organs were measured by counting with a gamma-counter. Radioactivity in the organs of the IP_3 -treated animals was compared with that of control animals which had been treated with saline for the same period of time. In the results radioactivity
30 in the organs of the animals treated with IP_3 is expressed as % of the radioactivity found in controls. The results were as follows:

35 Organ levels of mice treated with cadmium and IP_3 as per cent of control levels (controls = 100). 15 mice in each group. Said control group had been treated with saline and Cd as mentioned.

0179441

	<u>Organ</u>	<u>IP₃</u>
	Lung	80
	Heart	77
5	Aorta	89
	Spleen	81
	Salivary gland	82
10	Liver	100
	Kidney	102

The results show that IP₃ caused a reduction in cadmium
15 levels in all studied organs except liver and kidney at
which letter sites the Cd is believed to be relatively
safe.

20

25

30

35

Example 16

0179441

The effect of IP_3 on platelet aggregation after smoking in humans was studied.

5

Four young healthy male non-smokers received, on two occasions, a capsule containing 50 mg of IP_3 or 50 mg of a placebo. The IP_3 used was the Ca-salt of D-myo-inositol-1,2,6-triphosphate. Neither subject nor investigator knew whether the subject had received IP_3 or placebo.

Two hours after ingestion of the capsule, a blood sample was obtained. The subject then smoked two cigarettes in rapid succession. A second blood sample was obtained after smoking. The aggregation responses of the platelets to ADP and collagen in the two samples were determined, using essentially the same procedure as in Example 14. The results are expressed as change in aggregation from the pre-smoking to the post-smoking sample. A positive sign indicates that aggregation was stronger after smoking.

25	Aggregation agent	Concentration of aggregating agent	IP_3	Placebo	Difference between IP_3 and placebo
	ADP	0.5 mmol	+ 1.5	+ 7.25	5.85
	"	1 mmol	- 1.5	+ 0.25	1.75
	"	2.5 mmol	- 1.5	0	1.5
	"	5 mmol	- 2.5	- 0.75	1.75
30	Collagen	0.5 mg	+ 5.75	+ 12.25	6.5
	"	1 mg	- 8.25	+ 1.75	10.0
	"	2.5 mg	- 3.75	0	3.75
	"	5 mg	- 1.5	- 0.25	1.25

35 In the placebo group, smoking caused an increase in aggregation, which was most marked at low concentrations of aggregation agents. In all cases this effect was counteracted by IP_3 . Thus IP_3 prevents increase of platelet aggregation caused by smoking.

0179441

5 A reaction mixture consisting of 48 mmol KH_2PO_4 , 2 mmol Na-ascorbate, 0.1 mmol H_2O_2 , 0.5 mmol Fe and 1.7 mmol deoxyribose was incubated at 37°C for 1 hour. Similar reactions mixtures including EDTA 1 mmol or inositol-tri-phosphate (IP_3) 1 mmol were similarly incubated. The IP_3 used was D-myo-inositol-1.2.6- triphosphate.

10 After incubation 1.65 ml thiobarbituric acid in 50 mmol NaOH and 1.65 ml 2.8 % trichloroacetic acid was added to 2 ml of the reaction mixture. The mixture was heated to 100°C for 20 minutes and the absorbance at 532 nm
15 was measured with water as a blank.

The experiments were performed with iron in the form of Fe^{2+} ($\text{Fe}(\text{NH}_4)\text{SO}_4$) and Fe^{3+} (Fe Cl_3). The results were as follows:

20 Production of free radicals catalyzed by Fe^{2+} and Fe^{3+} in the presence of IP_3 or EDTA, expressed as absorbance at 532 nm.

25

<u>Group</u>	<u>Fe^{2+}</u>	<u>Fe^{3+}</u>
Control	0.76	0.79
EDTA	2.2	1.86
30 IP_3	0.46	0.43

35 These results show that the formation of free radicals in the reaction mixture was diminished by 40 % after addition of IP_3 . The addition of EDTA had an opposite effect. It strongly increased production of free radicals. Thus IP_3 was shown to reduce iron-dependent formation of free radicals.

Lipid peroxidation was studied in lipid micelles. The following reaction mixture was incubated for 2 hours at 37°C:

0.4 ml Clark-Lubs buffer pH 5.5
 0.2 ml phospholipid liposomes
 0.1 ml IP_3 0.5-5 mM or 0.1 ml H_2O
 0.1 ml Fe^{2+} 1 mM or 0.1 ml H_2O
 0.1 ml Al^{3+} 4 mM or 0.1 ml H_2O
 0.1 ml H_2O

The IP_3 was D-myo-inositol-1,2,6-triphosphate. After incubation, 0.5 ml of thiobarbituric acid + 0.5 ml 25 % HCl was added and the mixture was heated at 100°C for 15 minutes. 1 ml Lubrol PX 1% (Sigma) was added and lipid peroxidation was measured by measuring the absorbance at 532 nm. The results were as follows:

<u>Concentration, mM</u>				
<u>Experiment</u>	<u>Fe^{2+}</u>	<u>Al^{3+}</u>	<u>IP_3</u>	<u>Absorbance 532 nm</u>
1	0.1	0	0	0.367
2	0	0.4	0	0.128
3	0.1	0.4	0	0.896
4	0.1	0.4	0.5	0.367
5	0.1	0	0.5	0.303
6	0.1	0	0.4	0.260
7	0.1	0	0.2	0.297
8	0.1	0	0.1	0.283
9	0.1	0	0.05	0.271
10	0	0	0	0.133

Fe^{2+} caused lipid peroxidation (group 1 vs 10). Al^{3+} itself caused no peroxidation (2 vs 10) whereas the combination of Fe^{2+} + Al^{3+} caused much stronger peroxidation than Fe^{2+} alone (1 vs 3). Addition of IP_3 completely prevented the interaction between Fe^{2+} and Al^{3+} (3 vs 4). In systems with only Fe^{2+} , IP_3 caused marked reduction in radical formation (1 vs 5-9).

Example 19

Hydrolysis of phytic acid in wheat, extraction and analysis of IP_3 .

Ground wheat seeds, 100 g containing 1 % myo-inositol-hexaphosphate IP_6 was incubated in 1000 ml sodiumacetate buffer at pH 5.2 at 35°C. After an incubation period of 30 minutes, the slurry was frozen to -10°C in order to stop the hydrolysis.

15 10 g of the frozen material was extracted with 100 ml 0.4 M HCl. The suspension was shaken for 1 hr and subsequently centrifuged. The supernatant was collected and neutralized to pH 7 with an aqueous solution of NaOH. A sample of the supernatant was analyzed with HPLC. The analysis method was calibrated with welldefined inositol-phosphates. The IP_3 content of the extract was 10 mg inositoltriphosphate.

25

30

35

Example 20

Hydrolysis of phytic acid in white beans, extraction
5 and analysis of IP_3 .

The same method was used as described in Example 19
except for the difference that 100 g white beans contain-
10 ing 1 % myo-inositol hexaphosphate was incubated at
55°C for 10 hrs.

10 g of the frozen material was extracted with 100 ml
15 0.4 M HCl. The suspension was shaken for 1 hour and sub-
sequently centrifuged. The supernatant was collected
and neutralized to pH 7 with an aqueous solution of NaOH.
A sample of the supernatant was analyzed with HPLC. The
 IP_3 content of the extract was 5 mg inositoltriphosphate.

20

25

30

35

Example 21

Hydrolysis of phytic acid in soybeans after addition of a phytase
5 source from microorganisms, extraction and analysis of IP_3 .

300 g soy beans were soaked over night (1.4 % IP_6 content),
peeled and then boiled for 30 minutes. 3 ml water containing
10 about 1 g *Rhizopus oligosporus*, NRRL 2710 was added and
the mixture was incubated at 40°C for 20 hours. 10 g
of the mixture was extracted and analyzed by HPLC as
described in Example 19. The IP_3 content of the extract
was 160 mg.

15

20

25

30

35

Example 22

Hydrolysis of phytic acid in white beans with crude wheat
5 phytase, extraction and analysis of IP_3 .

Ground beans, 100 g, containing 1 % myo-inositol-hexa-
phosphate were suspended in 1000 ml sodiumacetate buffer
10 at pH 5.2. 500 mg crude wheat phytase (from Sigma Chemical
Co) was added. The mixture was incubated at $55^{\circ}C$ at shaking.
After an incubation period of 12 hrs the slurry was frozen
to $-10^{\circ}C$ in order to stop the hydrolysis.

15 10 g of the frozen material was extracted with 100 ml
0.4 M HCl. The suspension was shaken for 1 hour and sub-
sequently centrifuged. The supernatant was collected
and neutralized to pH 7 with an aqueous solution of NaOH.
20 A sample of the supernatant was analyzed with HPLC. The
 IP_3 content of the extract was 40 mg IP_3 .

25

30

35

Example 23

Content of IP_3 in white beans after addition of sodium-
5 phytate and hydrolysis.

0.3 gram of sodiumphytate was added to 100 g ground white
beans (1 % IP_6). The mixture was incubated in 1000 ml
10 sodiumacetate buffer at pH 5.2 at 55°C. After an incubation
period of 4 hours the slurry was frozen to -10°C in
order to stop the hydrolysis. 10 g of the frozen material
was extracted and analyzed by HPLC as described in Example
19. The IP_3 content of the extract was 15 mg IP_3 .

15

20

25

30

35

Example 24

1.0 kg of rice bran, containing ca 1 % inositolhexaphosphate
5 (IP₆) was suspended in 10 l sodiumacetate buffer at pH
5 at 25°C. After 4 hours when 50 % inorganic phosphorus
had been released the slurry was extracted with an ad-
dition of 1 l 2 M HCl. The suspension was shaken for
1 hour and subsequently centrifuged. The supernatant
10 was neutralized to pH 7 with an aqueous solution of Ca(OH)₂.
A precipitate was obtained when 5 l ethanol was added.
The calciumsalt consisting of a composition of different
inositolphosphates was centrifuged, dried and recrysta-
lized. 20 mg of the recrystallized calciumsalt was con-
15 verted to the acid form by addition of diluted hydrochloric
acid and was analyzed by HPLC. The composition consisted
of 20 % inositoltriphosphate. The rest consisted of other
inositolphosphates.

20

25

30

35

0179441

Example 25

Hydrolysis of sodium phytate with wheat phytase and
fractionation of a mixture of inositolphosphates.

5

A 1.6 gram quantity of sodium phytate (from corn, Sigma
Chemical Co) was dissolved in 650 ml sodium acetate buffer,
pH 5.2. 2.7 gram wheat phytase (EC 3.1.3.26, 0.015 U/mg,
from Sigma Chemical Co) was added and the mixture was
incubated at 38°C.

10

The dephosphorylation was followed by determining the
inorganic phosphorus released. After 3 hours when 50%
inorganic phosphorus was liberated the hydrolysis was
stopped by adding 30 ml ammonia to pH 12. A liquid mixture
containing inositolphosphates was obtained.

15

350 ml of the mixture was passed through an ion-exchange
column (Dowex 1, chloride form, 25 mm x 250 mm) and eluted
with a linear gradient of hydrochloric acid (0-0.7 N HCl).
Aliquots of eluted fractions were completely hydrolyzed
in order to determine the contents of phosphorus and
inositol. The peaks correspond to different inositol-
phosphates i.e. a peak with the ratio of phosphorus to
inositol of three to one consists of inositoltriphosphate
etc. Two fractions with the ratio of phosphorus to inositol
of three to one were obtained.

25

30

35

Example 26

Fractionation of inositoltriphosphates.

5

100 ml of the first fraction obtained in Example 25 with a phosphorus/inositol ratio of three to one was neutralized and precipitated as a barium salt after addition of 10 % excess of 0.1 M barium acetate solution. 600 mg of the precipitated salt was dissolved in 50 ml diluted hydrochloric acid. The solution was separated on an ion-exchange column (Dowex 1, chloride form, 25 mm x 2500 mm) with diluted hydrochloric acid as eluent. Aliquots of eluted fractions were analyzed for phosphorus. Three peaks consisting of isomers of inositoltriphosphates can be seen.

20

25

30

35

Example 27

5 Structural determination of isomers of inositol-triphosphates
with NMR.

10 The three peaks obtained in Example 26 were analyzed
by H-NMR. Data show that the peaks consist of myo-inositol-
1.2.6-triphosphate, myo-inositol-1.2.3-triphosphate and
myo-inositol-1.3.4-triphosphate respectively.

15 The second fraction obtained in Example 25 with a phos-
phorus/inositol ratio of three to one was analyzed by
H-NMR. Data show that the fraction consists of myo-inositol-
1.2.5-triphosphate.

20

25

30

35

0179441

Example 28

Determination of optical isomers of inositol-triphosphates.

5 20 mg of the compounds determined with NMR according
to Example 27 to be myo-inositol-1.2.6-triphosphate and
myo- inositol-1.3.4-triphosphate were further chromato-
graphed on a chiral column based on acetylated cellulose
10 (20 mm x 300 mm from Merck) with a mixture of ethanol
and water as eluent. The fractions were analyzed with
a polarimeter. As can be seen each compound consists
of one optical isomer, D-myo-inositol-1.2.6-triphosphate
and L-myo-inositol-1.3.4-triphosphate respectively.

15

20

25

30

35

Example 29

Hydrolysis of sodium phytate with baker's yeast and fractionation of a mixture of inositolphosphates.

5


A 0.7 gram quantity of sodium phytate (from corn, Sigma Chemical Co) was dissolved in 600 ml sodium acetate buffer pH 4.6. 50 gram of baker's yeast from Jästbolaget, Sweden (dry substance: 28 %, nitrogen content: 2 %; phosphorus content: 0.4 %) was added with stirring and incubation was continued at 45°C. The dephosphorylation was followed by determining the inorganic phosphorus released. After 7 hours when 50% inorganic phosphorus was liberated the hydrolysis was stopped by adding 30 ml of ammonia to pH 12. The suspension was centrifuged and the supernatant was collected.

400 ml of the supernatant was passed through an ion-exchange column (Dowex 1, chloride form, 25 mm x 250 mm) and eluted with a linear gradient of hydrochloric acid (0-0.7 N HCl).

Aliquots of eluted fractions were completely hydrolyzed in order to determine the contents of phosphorus and inositol. The peaks correspond to different inositol-phosphates i.e. a peak with the ratio of phosphorus to inositol of three to one consists of inositoltriphosphates etc.


35

0179441



D-myo-inositol-1,2,5-tri-
phosphate alternatively
D-1,2,5-IP₃


5



myo-inositol-1,2,3-tri-
phosphate alternatively
1,2,3-IP₃


10

P = O-PO₃H₂




L-myo-inositol-1,3,4-tri-
phosphate alternatively
L-1,3,4-IP₃

15




L-myo-inositol-1,2-diphos-
phate alternatively L-1,2-IP₂

20



D-myo-inositol-1,2,5,6-
tetra-phosphate or D-1,2,5,6-
IP₄

25



L-myo-inositol-1,2,3,4,5-
penta phosphate or
L-1,2,3,4,5-IP₅

30

P = -I-PO₃H₂

0179441

Claims

1. A method of making a food composition, said food being initially substantially free of inositoltriphosphate (IP_3), which comprises adding a source of IP_3 to the composition in an amount sufficient to provide a final
5 concentration of at least 5 mg IP_3 per 100 g of composition.
2. A method according to claim 1, wherein said source is IP_3 .
- 10 3. A method according to claim 1, wherein said source is inositolhexaphosphate (IP_6), inositolpentaphosphate (IP_5) and/or inositoltetraphosphate (IP_4) in the presence of phytase.
- 15 4. A method according to claim 1, wherein the source is at least one of the following IP_3 isomers, D-myo-inositol-1.2.6-triphosphate, D-myo-inositol-1.2.5-triphosphate, myo-inositol-1.2.3-triphosphate, L-myo-inositol-1.3.4-triphosphate or D-myo-inositol-1.4.5-triphosphate.
- 20 5. A method according to claim 1, wherein in the materials composing the composition, a content of phytase and of an inositolphosphate selected from the group consisting of IP_4 , IP_5 and IP_6 is established and at at least one
25 stage of the production process, the time and the temperature of the processing as well as the pH-value are controlled to allow an incubation in such a way that a content of IP_3 of at least 20 mg per 100 g composition is obtained.
- 30 6. A method according to claim 1, wherein the composition is solid and the content of IP_3 is at least 20 mg per 100 g dry composition.

7. A method according to claim 1, wherein the composition is a cereal based material.

0179441

5 8. A method according to claim 7, wherein the composition is selected from the group consisting of breakfast cereals, cakes and biscuits.

10 9. A method according to claim 1, wherein the composition is selected from the group consisting of sweets, chocolates and chewing gums.

10. A method according to claim 1, wherein the composition is a vegetable.

15 11. A method according to claim 6, wherein the composition is bread and the content of IP_3 is at least 50 mg per 100 g dry bread.

20 12. A method according to claim 1, wherein the composition is selected from the group consisting of beverages and soups.

25 13. A method according to claim 5, wherein the incubation is carried out separately and the IP_3 containing product obtained is added to the composition to control the desired IP_3 content.

14. A method according to claim 3, wherein the phytase is added in the form of yeast.

30 15. A method according to claim 5, wherein the temperature ranges between 20 and 70°C and the pH-value is between 4 and 8 at the incubation step.

35 16. A method according to claim 1, wherein the IP_3 concentration is between 20 and 500 mg per 100 g dry composition.

17. A method according to claim 1, wherein the IP_3 concentration is between 50 and 500 mg per 100 g dry composition.

0179441

18. A method according to claim 1, wherein the composition is in liquid form and the IP_3 concentration is between 5 and 300 mg per 100 g of the liquid composition.
19. A method of making a food composition containing initially less than about 10 mg of IP_3 per 100 g of composition wherein the content of IP_3 is increased to at least 20 mg per 100 g of composition by addition of IP_3 or a source thereof or conversion of initially contained inositolphosphate selected from the group consisting of IP_6 , IP_5 and IP_4 by enzymatic processing.
20. A method according to claim 19, wherein the composition is selected from the group consisting of cereal based material, sweets, chocolates, chewing gums, vegetables, fruits, beverages and soups.
21. A method according to claim 19, wherein the content of IP_3 is increased to at least 30 mg and is preferably in the range of 50-500 mg per 100 g composition.
22. A method of making a food composition, wherein in the materials composing the composition, a content of phytase and of an inositolphosphate selected from the group consisting of IP_4 , IP_5 and IP_6 is established and at at least one stage of the production process, the time and the temperature of the processing as well as the pH-value are controlled to allow an incubation in such way that the degree of hydrolysis of IP_4 , IP_5 and IP_6 is between 40 and 70% to obtain a sufficient content of IP_3 .
23. A non cereal based food composition having a content of IP_3 of at least 5 mg per 100 g of the composition.
24. A composition according to claim 23 wherein the content of IP_3 is at least 10 mg and preferably in the range of 20-500 mg per 100 g of composition.

0179441

25. A composition according to claim 23, wherein the composition is selected from the group consisting of sweets, chocolates, chewing gums, vegetables, fruits, beverages and soups.
- 5 26. A composition according to claim 23, wherein the IP_3 is at least one of the following isomers, D-myo-inositol-1.2.6-triphosphate, D-myo-inositol-1.2.5-triphosphate, myo-inositol-1.2.3-triphosphate, L-myo-inositol-1.3.4-triphosphate or D-myo-inositol-1.4.5-triphosphate.
- 10 27. A bakery food composition, wherein the concentration of IP_3 is at least 20 mg per 100 g of dry composition, wherein the degree of hydrolysis of naturally contained IP_6 and/or of added IP_6 is between 40 and 70% .
- 15 28. A composition according to claim 27, wherein the concentration of IP_3 is at least 30 mg and preferably in the range of 50-500 mg per 100 g of dry composition.
- 20 29. A composition according to claim 27 wherein the bakery food is selected from the group consisting of bread, breakfast cereals, bisquits and cakes.
- 25 30. A composition according to claim 27, wherein the IP_3 is substantially D-myo-inositol-1.2.6-triphosphate.
- 30 31. A composition according to any of claims 23-27, wherein the IP_3 is a hydrolysate product derived from an inositol-phosphate selected from IP_4 , IP_5 and IP_6 in which the distribution curve showing the content of the different inositolphosphates has a maximum for IP_3 .
- 35 32. A composition according to claim 31, wherein the content of IP_3 is more than 40% by weight of the total amount of inositolphosphates and 30 to 85% by weight of the remaining inositolphosphates consist of IP_2 and IP_4 .

0179441

33. A composition according to claim 23, in concentrate form which comprises IP_3 in a food acceptable extending agent.

5 34. A food composition, wherein the concentration of IP_3 is at least 20 mg per 100 g of dry composition, and IP_3 has been added to the composition and/or formed by adding phytase and an inositolphosphate selected from the group consisting of IP_4 , IP_5 and IP_6 .

10

15

20

25

30

35

THIS PAGE BLANK (USPTO)